

Turf out of Turf Guide

Undesired Turf		Desired Turf	Product Recommendation and Notes*
Bermudagrass	Out of	Bahiagrass	—
Bermudagrass	Out of	St. Augustine	Prograss 1.5L (2 gal/acre) or 4SC (3 qt/acre) + Atrazine 4L (2 gal/acre) Spring apps should start in mid March. Repeat in 30 days.
Bermudagrass	Out of	Seashore Paspalum	Prograss + Cutless (1.5 + 0.75 lb a.i./acre)
Bermudagrass	Out of	Zoysiagrass	Fusilade T&O II 2L (5 oz/acre) + Turflon Ester 4L (1 qt/acre) Add a nonionic surfactant. Begin treatment on zoysiagrass at 3–4 oz/A in early June. Adding triclopyr ester increases control but should not be used on warm-season grasses unless temporary phytotoxicity is acceptable.
Bermudagrass	Out of	Zoysiagrass (Meyer)	Acclaim Extra 0.94L (24 oz/acre) + Turflon Ester 4L (1 qt/acre) Apply in late spring or early summer to actively growing plants. Do not apply to moisture- or heat- stressed turf or weeds. Repeat in 2 -3 weeks for complete control. Adding triclopyr ester increases control but should not be used on warm-season grasses unless temporary phytotoxicity is acceptable.
Bermudagrass	Out of	Bentgrass Fairways	Acclaim Extra 0.94L (24 oz/acre) + Turflon Ester 4L (1 qt/acre) Apply in late spring or early summer to actively growing plants. Do not apply to moisture- or heat- stressed turf or weeds. Repeat in 2 -3 weeks for complete control.
Bermudagrass Suppression	Out of	Creeping Bentgrass	Tupersan 50WP (siduron) (1lb/1,000 sq ft) Or Acclaim Extra 0.57L + Turflon Ester 4L (39 oz + 1 pt/acre) Granular formulations of siduron also are available. Control is generally best with spring (March + April + May + early June) and fall (late September + October + November) applications when the bentgrass is actively growing and the bermudagrass is not. Acclaim is for fairway use only.
Bermudagrass	Out of	Ryegrass	Acclaim Extra 0.94L (24 oz/acre) + Turflon Ester 4L (1 qt/acre) Apply in late spring or early summer to actively growing plants. Do not apply to moisture- or heat- stressed turf or weeds. Repeat in 2 -3 weeks for complete control.
Bermudagrass	Out of	Poa Annua	—
Bermudagrass	Out of	Kentucky Bluegrass	Acclaim Extra 0.94L (24 oz/acre) + Turflon Ester 4L (1 qt/acre) Apply in late spring or early summer to actively growing plants. Do not apply to moisture- or heat- stressed turf or weeds. Repeat in 2 -3 weeks for complete control.
Bermudagrass	Out of	Tall Fescue	Fusilade T&O II 2L (5 oz/acre) + Turflon Ester 4L (1 qt/acre) Add a nonionic surfactant. Make first app in spring after bermudagrass green-up and a second application in early fall. Or Acclaim Extra 0.94L (24 oz/acre) + Turflon Ester 4L (1 qt/acre) Apply in late spring or early summer to actively growing plants. Do not apply to moisture- or heat- stressed turf or weeds. Repeat in 2 -3 weeks for complete control.

*Provided by L.B. McCarty, Clemson University. — Indicates no recommendations.

Please contact your
Harrell's Representative
for more information



720 Kraft Road
Lakeland, FL 33815
800.282.8007
www.harrells.com

Turf out of Turf Guide

Undesired Turf		Desired Turf	Product Recommendation and Notes*
Bermudagrass	Out of	Fine Fescue	Vantage 1L or Segment 1L (2.25 pt/acre) Apply before weeds mature. Or Acclaim Extra 0.94L (24 oz/acre) + Turflon Ester 4L (1 qt/acre) Apply in late spring or early summer to actively growing plants. Do not apply to moisture- or heat- stressed turf or weeds. Repeat in 2-3 weeks for complete control.
Bahiagrass	Out of	Bermudagrass	Manor or Blade 60DF (metsulfuron) (0.25-1 oz/acre) Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.
Bahiagrass	Out of	St. Augustine	Manor or Blade 60DF (metsulfuron) (0.25-1 oz/acre) Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.
Bahiagrass	Out of	Seashore Paspalum	—
Bahiagrass	Out of	Zoysiagrass (Meyer & Emerald)	Manor or Blade 60DF (metsulfuron) (0.25-1 oz/acre) Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.
Bahiagrass	Out of	Bentgrass	—
Bahiagrass	Out of	Ryegrass	—
Bahiagrass	Out of	Poa Annua	—
Bahiagrass	Out of	Kentucky Bluegrass	Manor or Blade 60DF (metsulfuron) (0.25-1 oz/acre) Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.
Bahiagrass	Out of	Fine Fescue	Manor or Blade 60DF (metsulfuron) (0.25-1 oz/acre) Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.
St Augustine	Out of	Bahiagrass	—
St. Augustine	Out of	Bermudagrass	MSMA (1.8 pt/acre) Repeat applications 7 days apart as needed. Or Drive 75 (1 lb/acre)
St. Augustine	Out of	Seashore Paspalum	—

*Provided by L.B. McCarty, Clemson University. — Indicates no recommendations.

Please contact your
Harrell's Representative
for more information



720 Kraft Road
Lakeland, FL 33815
800.282.8007
www.harrells.com

Turf out of Turf Guide

Undesired Turf		Desired Turf	Product Recommendation and Notes*
St. Augustine	Out of	Zoysiagrass	MSMA (1.8 pt/acre) Repeat applications 7 days apart as needed. Or Drive 75 (1 lb/acre)
St. Augustine	Out of	Zoysiagrass (Meyer)	Acclaim Extra 0.94L (24 oz/acre) + Turflon Ester 4L (1 qt/acre) Apply in late spring or early summer to actively growing plants. Do not apply to moisture- or heat- stressed turf or weeds. Repeat in 2 -3 weeks for complete control. The addition of triclopyr ester increased control but should not be used on warm-season grasses unless temporary phytotoxicity is acceptable.
St. Augustine	Out of	Bentgrass Fairways	Acclaim Extra 0.94L (24 oz/acre) + Turflon Ester 4L (1 qt/acre) Apply in late spring or early summer to actively growing plants. Do not apply to moisture- or heat- stressed turf or weeds. Repeat in 2 -3 weeks for complete control.
St. Augustine	Out of	Ryegrass	Acclaim Extra 0.94L (24 oz/acre) + Turflon Ester 4L (1 qt/acre) Apply in late spring or early summer to actively growing plants. Do not apply to moisture- or heat- stressed turf or weeds. Repeat in 2 -3 weeks for complete control.
St. Augustine	Out of	Poa Annua	—
St. Augustine	Out of	Kentucky Bluegrass	Acclaim Extra 0.94L (24 oz/acre) + Turflon Ester 4L (1 qt/acre) Apply in late spring or early summer to actively growing plants. Do not apply to moisture- or heat- stressed turf or weeds. Repeat in 2 -3 weeks for complete control.
St. Augustine	Out of	Fine Fescue	Acclaim Extra 0.94L (24 oz/acre) + Turflon Ester 4L (1 qt/acre) Apply in late spring or early summer to actively growing plants. Do not apply to moisture- or heat- stressed turf or weeds. Repeat in 2 -3 weeks for complete control.
St. Augustine	Out of	Tall Fescue	Acclaim Extra 0.94L (24 oz/acre) + Turflon Ester 4L (1 qt/acre) Apply in late spring or early summer to actively growing plants. Do not apply to moisture- or heat- stressed turf or weeds. Repeat in 2 -3 weeks for complete control.
Seashore Paspalum	Out of	Bermudagrass	MSMA (1.8 pt/acre) Or TranXit GTA 25DG (2 oz/acre) Or Monument 75WG (0.56 oz/acre) Repeat applications 7 days apart are needed for control. Tank-mixing Sencor 25DG at 0.33 lb/acre with MSMA increases control as well as phyto potential. Add MSO to Monument or TranXit for increased activity.
Seashore Paspalum	Out of	St. Augustine	—
Seashore Paspalum	Out of	Bahiagrass	—

*Provided by L.B. McCarty, Clemson University. — Indicates no recommendations.

Please contact your
Harrell's Representative
for more information



720 Kraft Road
Lakeland, FL 33815
800.282.8007
www.harrells.com

Turf out of Turf Guide

Undesired Turf		Desired Turf	Product Recommendation and Notes*
Seashore Paspalum	Out of	Zoysiagrass	MSMA (1.8 pt/acre) Or TranXit GTA 25DG (2 oz/acre) Or Monument 75WG (0.56 oz/acre) Repeat applications 7 days apart are needed for control. Tank-mixing Sencor 25DG at 0.33 lb/acre with MSMA increases control as well as phyto potential. Add MSO to Monument or TranXit for increased activity.
Seashore Paspalum	Out of	Bentgrass	—
Seashore Paspalum	Out of	Ryegrass	—
Seashore Paspalum	Out of	Poa Annua	—
Seashore Paspalum	Out of	Kentucky Bluegrass	—
Seashore Paspalum	Out of	Fescue	—
Zoysiagrass	Out of	Bahiagrass	—
Zoysiagrass	Out of	St. Augustine	—
Zoysiagrass	Out of	Seashore Paspalum	—
Zoysiagrass	Out of	Bermudagrass	—
Zoysiagrass	Out of	Bentgrass	—
Zoysiagrass	Out of	Ryegrass	—
Zoysiagrass	Out of	Poa Annua	—
Zoysiagrass	Out of	Kentucky Bluegrass	—
Zoysiagrass	Out of	Fescue	—

*Provided by L.B. McCarty, Clemson University. — Indicates no recommendations.

Please contact your
Harrell's Representative
for more information



720 Kraft Road
Lakeland, FL 33815
800.282.8007
www.harrells.com

Turf out of Turf Guide

Undesired Turf		Desired Turf	Product Recommendation and Notes*
Bentgrass	Out of	Bermudagrass	Atrazine or Simazine (1 qt/acre) Or Kerb 50W (2 lb/acre) or TranXit (0.5-2 oz/acre) Or Monument (0.11-0.33 oz/acre) Or Revolver 0.19L (9-27 oz/acre) Use only on bermudagrass or possibly zoysiagrass. Do not apply on or up-slope to desirable bentgrass or overseeded turf as these may run. Movement is encourage when saturated soils are treated and/or heavy (>0.25 in) rainfall occurs within 48 hours of application. Time required for control increases as weeds mature, therefore apply in late fall for optimum results. For slow (3 to 6 weeks) transition, use the low rate of each herbicide listed. For quick transition (1 to 2 weeks), use TranXit, Revolver, or Monument at the high rate in mid-May. Treated plants do not show herbicide symptoms until air temperatures are consistently above 60F. Revolver is used in late summer or early fall to provide suppression of dallisgrass when used in spot treatments. Use Revolver at 1.5 – 2.0 fl oz per gallon and spray to wet followed 7 to 14 days later with an additional spot treatment. If regrowth occurs, a third application may be necessary.
Bentgrass	Out of	St. Augustine	Atrazine or Simazine (1 qt/acre) Best results are late summer or early fall applications. Or Kerb 50W (2 lb/acre)
Bentgrass	Out of	Seashore Paspalum	—
Bentgrass	Out of	Zoysiagrass	—
Bentgrass	Out of	Bahiagrass	—
Bentgrass	Out of	Ryegrass	—
Bentgrass	Out of	Poa Annua	—
Bentgrass	Out of	Kentucky Bluegrass	Tenacity 4L (4-8 oz/acre) Treat twice, 3 weeks apart.
Bentgrass	Out of	Fine Fescue	Tenacity 4L (4-8 oz/acre) Treat twice, 3 weeks apart.
Bentgrass	Out of	Tall Fescue	Tenacity 4L (4-8 oz/acre) Treat twice, 3 weeks apart.
Bentgrass	Out of	St. Augustine Sod	Tenacity 4L (4-8 oz/acre) Treat twice, 3 weeks apart. Use low rate.
Ryegrass	Out of	Bahiagrass	—

*Provided by L.B. McCarty, Clemson University. — Indicates no recommendations.

Please contact your
Harrell's Representative
for more information



720 Kraft Road
Lakeland, FL 33815
800.282.8007
www.harrells.com

Turf out of Turf Guide

Undesired Turf		Desired Turf	Product Recommendation and Notes*
Ryegrass	Out of	St. Augustine	Atrazine or Simazine (1 qt/acre) Or Kerb 50W (2 lb/acre) Or Manor or Blade 60DF (metsulfuron) 0.25-1 oz/acre Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.
Ryegrass	Out of	Seashore Paspalum	—
Ryegrass	Out of	Zoysiagrass (Meyer & Emerald)	Manor or Blade 60DF (metsulfuron) 0.25-1 oz/acre Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.
Ryegrass	Out of	Bermudagrass	Atrazine or Simazine (1 qt/acre) Or Kerb 50W (2 lb/acre) Or Manor or Blade 60DF (metsulfuron) (0.25-1 oz/acre) Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible. Or TranXit (0.5-2 oz/acre) Or Monument (0.11-0.33 oz/acre) Or Revolver 0.19L (9-27 oz/acre) Use only on bermudagrass or possibly zoysiagrass. Do not apply on or up-slope to desirable bentgrass or overseeded turf as these may run. Movement is encourage when saturated soils are treated and/or heavy (>0.25 in) rainfall occurs within 48 hours of application. Time required for control increases as weeds mature, therefore apply in late fall for optimum results. For slow (3 to 6 weeks) transition, use the low rate of each herbicide listed. For quick transition (1 to 2 weeks), use TranXit, Revolver, or Monument at the high rate in mid-May. Treated plants do not show herbicide symptoms until air temperatures are consistently above 60F. Revolver is used in late summer or early fall to provide suppression of dallisgrass when used in spot treatments. Use Revolver at 1.5 – 2.0 fl oz per gallon and spray to wet followed 7 to 14 days later with an additional spot treatment. If regrowth occurs, a third application may be necessary.
Ryegrass	Out of	Bentgrass	—
Ryegrass	Out of	Poa Annua	—

*Provided by L.B. McCarty, Clemson University. — Indicates no recommendations.

Please contact your
Harrell's Representative
for more information



720 Kraft Road
Lakeland, FL 33815
800.282.8007
www.harrells.com

Turf out of Turf Guide

Undesired Turf		Desired Turf	Product Recommendation and Notes*
Ryegrass	Out of	Kentucky Bluegrass	Manor or Blade 60DF (metsulfuron) 0.25-1 oz/acre Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.
Ryegrass	Out of	Fine Fescue	Manor or Blade 60DF (metsulfuron) 0.25-1 oz/acre Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.
Perennial Ryegrass	Out of	Bahia	Corsair 75DF up to 5.3 oz/acre
Perennial Ryegrass	Out of	Bermudagrass	Corsair 75DF up to 5.3 oz/acre Or Kerb 50W (2-3 lb/acre) Do not apply on or up-slope to desirable Bentgrass or overseeded turf it may run.
Perennial Ryegrass	Out of	Bentgrass Fairways	Corsair 75DF up to 5.3 oz/acre. Spot treat using a hand sprayer delivering 1 gallon of spray per 1000 sq. ft.
Perennial Ryegrass	Out of	Fine Fescue	Corsair 75DF up to 5.3 oz/acre. Spot treat using a hand sprayer delivering 1 gallon of spray per 1000 sq. ft.
Perennial Ryegrass	Out of	Kentucky Bluegrass	Corsair 75DF up to 5.3 oz/acre. Spot treat using a hand sprayer delivering 1 gallon of spray per 1000 sq. ft.
Poa Annua	Out of	Bermudagrass	Kerb 50W (2-3 lb/acre) Do not apply on or up-slpoe to desirable bengrass or overseeded turf. It may run.
Poa Annua	Out of	St. Augustine	—
Poa Annua	Out of	Seashore Paspalum	—
Poa Annua	Out of	Zoysiagrass	—
Poa Annua	Out of	Bahiagrass	—
Poa Annua	Out of	Perennial Ryegrass Fairways	Velocity 17.6 SC (6-12 oz/acre) Apply between Feb. 1 and March 15 when daytime/nighttime temperatures are 70/50 F at 6 to 12 oz/acre in 25 to 50 gallons of water. Reapply in 21 days. Use higher labeled rates as Poa matures. Higher rates, however, may cause short-term ryegrass chlorosis. Treated ryegrass should be overseeded before Oct. 15th at >300 lbs seed/acre. No surfactant or adjuvants are needed. On bentgrass fairways, apply 2 to 6 oz/acre weekly in spring/summer only when the bentgrass is actively growing. For Poa trivialis control, apply 4 to 9 oz/acre weekly. For all scenarios, use lower rates when higher weed populations are present to prevent voids from developing.

*Provided by L.B. McCarty, Clemson University. — Indicates no recommendations.

Please contact your
Harrell's Representative
for more information



720 Kraft Road
Lakeland, FL 33815
800.282.8007
www.harrells.com

Turf out of Turf Guide

Undesired Turf		Desired Turf	Product Recommendation and Notes*
Poa Annua	Out of	Creeping Bentgrass Fairways	Velocity 17.6 SC (6-12 oz/acre) Apply between Feb. 1 and March 15 when daytime/nighttime temperatures are 70/50 F at 6 to 12 oz/acre in 25 to 50 gallons of water. Reapply in 21 days. Use higher labeled rates as Poa matures. Higher rates, however, may cause short-term ryegrass chlorosis. Treated ryegrass should be overseeded before Oct. 15th at >300 lbs seed/acre. No surfactant or adjuvants are needed. On bentgrass fairways, apply 2 to 6 oz/acre weekly in spring/summer only when the bentgrass is actively growing. For Poa trivialis control, apply 4 to 9 oz/acre weekly. For all scenarios, use lower rates when higher weed populations are present to prevent voids from developing. Or Prograss 1.5EC (3-4 pt/acre) or 4SC (1-1.5 pt/acre) Treat young (1 to 5 leaf stage) weeds in fall. Use lower rate on closer mowed turf. Will not adequately control mature plants or perennial biotypes. Multiple applications spaced 3 weeks apart may be necessary. Do not use on turf less than 8 weeks old nor reseed within 6 weeks after application. Bentgrass that is shaded, poorly drained (wet), and cold often experience herbicide damage.
Poa Annua	Out of	Creeping Bentgrass	Trimmit (paclobutrazol) (8-16 oz/acre) Apply 30 days apart at higher rate 3 times in fall (September to early Dec.) plus 2 or 3 times in very early spring (late Feb. to mid April) when bentgrass is actively growing. Increased Poa control often occurs at the lower rate if a sterol inhibitor fungicide (DMI) such as Banner Maxx at 1 oz/1000 sq.ft. is applied 2 weeks following each paclobutrazol applications. Do not use if Poa annua populations exceed 70% as severe stand thinning or discoloration may result. Do not apply within 4 weeks of anticipated cold or hot weather. Note: This program is designed as a gradual transition or conversion from Poa annua to bentgrass. Repeat applications over several years will be required. Treated Poa will appear noticeably lighter green in color while treated bentgrass may appear 'grainy.' Apply only to actively growing bentgrass
Poa Annua	Out of	Kentucky Bluegrass Fairways & Roughs	Prograss 1.5EC (3-4 pt/acre or 4SC (1-1.5 pt/acre) Treat young (1 to 5 leaf stage) weeds in fall. Use lower rate on closer mowed turf. Will not adequately control mature plants or perennial biotypes. Multiple applications spaced 3 weeks apart may be necessary. Do not use on turf less than 8 weeks old nor reseed within 6 weeks after application. Bentgrass that is shaded, poorly drained (wet), and cold often experience herbicide damage.
			—
Poa Annua	Out of	Tall Fescue	Prograss 1.5EC (3-4 pt/acre) or 4SC (1-1.5 pt/acre) Treat young (1 to 5 leaf stage) weeds in fall. Use lower rate on closer mowed turf. Will not adequately control mature plants or perennial biotypes. Multiple applications spaced 3 weeks apart may be necessary. Do not use on turf less than 8 weeks old nor reseed within 6 weeks after application. Bentgrass that is shaded, poorly drained (wet), and cold often experience herbicide damage.
Poa Trivialis	Out of	Perennial Ryegrass Fairways	Velocity 17.6 SC (6-12 oz/acre) Apply between Feb. 1 and March 15 when daytime/nighttime temperatures are 70/50 F at 6 to 12 oz/acre in 25 to 50 gallons of water. Reapply in 21 days. Use higher labeled rates as Poa matures. Higher rates, however, may cause short-term ryegrass chlorosis. Treated ryegrass should be overseeded before Oct. 15th at >300 lbs seed/acre. No surfactant or adjuvants are needed. On bentgrass fairways, apply 2 to 6 oz/acre weekly in spring/summer only when the bentgrass is actively growing. For Poa trivialis control, apply 4 to 9 oz/acre weekly. For all scenarios, use lower rates when higher weed populations are present to prevent voids from developing.
Poa Trivialis	Out of	Creeping Bentgrass Fairways	Velocity 17.6 SC (6-12 oz/acre) Apply between Feb. 1 and March 15 when daytime/nighttime temperatures are 70/50 F at 6 to 12 oz/acre in 25 to 50 gallons of water. Reapply in 21 days. Use higher labeled rates as Poa matures. Higher rates, however, may cause short-term ryegrass chlorosis. Treated ryegrass should be overseeded before Oct. 15th at >300 lbs seed/acre. No surfactant or adjuvants are needed. On bentgrass fairways, apply 2 to 6 oz/acre weekly in spring/summer only when the bentgrass is actively growing. For Poa trivialis control, apply 4 to 9 oz/acre weekly. For all scenarios, use lower rates when higher weed populations are present to prevent voids from developing. Or Certainty 75WDG (0.5 oz/acre) Start in spring when daytime temps. are >70 F (21 C). Three applications spaced 2 weeks apart are generally needed.

*Provided by L.B. McCarty, Clemson University. — Indicates no recommendations.

Please contact your
Harrell's Representative
for more information



720 Kraft Road
Lakeland, FL 33815
800.282.8007
www.harrells.com

Turf out of Turf Guide

Undesired Turf		Desired Turf	Product Recommendation and Notes*
Poa Trivialis	Out of	Bermudagrass	TranXit (0.5-2 oz/acre) Or Monument (0.11-0.33 oz/acre) Or Revolver 0.19L (9-27 oz/acre) Use only on bermudagrass or possibly zoysiagrass. Do not apply on or up-slope to desirable bentgrass or overseeded turf as these may run. Movement is encourage when saturated soils are treated and/or heavy (>0.25 in) rainfall occurs within 48 hours of application. Time required for control increases as weeds mature, therefore apply in late fall for optimum results. For slow (3 to 6 weeks) transition, use the low rate of each herbicide listed. For quick transition (1 to 2 weeks), use TranXit, Revolver, or Monument at the high rate in mid-May. Treated plants do not show herbicide symptoms until air temperatures are consistently above 60F. Revolver is used in late summer or early fall to provide suppression of dallisgrass when used in spot treatments. Use Revolver at 1.5 – 2.0 fl oz per gallon and spray to wet followed 7 to 14 days later with an additional spot treatment. If regrowth occurs, a third application may be necessary.
Kentucky Bluegrass	Out of	Bahiagrass	—
Kentucky Bluegrass	Out of	St. Augustine	Atrazine or Simazine (1 qt/acre) Or Kerb 50W (2 lb/acre)
Kentucky Bluegrass	Out of	Seashore Paspalum	—
Kentucky Bluegrass	Out of	Zoysiagrass	—
Kentucky Bluegrass	Out of	Bermudagrass	Atrazine or Simazine (1 qt/acre) Or Kerb 50W (2 lb/acre)
Kentucky Bluegrass	Out of	Ryegrass	—
Kentucky Bluegrass	Out of	Poa Annua	—
Kentucky Bluegrass	Out of	Bentgrass	—
Kentucky Bluegrass	Out of	Fescue	—
Fine Fescue	Out of	Bermudagrass	Atrazine or Simazine (1 qt/acre) Or Kerb 50W (2 lb/acre)
Fine Fescue	Out of	St. Augustine	Atrazine or Simazine (1 qt/acre) Or Kerb 50W (2 lb/acre)
Fescue	Out of	Seashore Paspalum	—
Fescue	Out of	Zoysiagrass	—
Fescue	Out of	Bahiagrass	—
Fescue	Out of	Ryegrass	—
Fescue	Out of	Poa Annua	—

*Provided by L.B. McCarty, Clemson University. — Indicates no recommendations.

Please contact your
Harrell's Representative
for more information



720 Kraft Road
Lakeland, FL 33815
800.282.8007
www.harrells.com

Turf out of Turf Guide

Undesired Turf		Desired Turf	Product Recommendation and Notes*
Tall Fescue	Out of	Kentucky Bluegrass	Corsair 75DF up to 5.3 oz/acre. Spot treat using a hand-help sprayer delivering 1 gal of spray solution per 1000 sq. ft. Spray to only wet the tall fescue blades. Or Manor or Blade 60DF (metsulfuron) 0.25-1 oz/acre Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.
Tall Fescue	Out of	Bentgrass Fairways	Corsair 75DF up to 5.3 oz/acre. Spot treat using a hand-help sprayer delivering 1 gal of spray solution per 1000 sq. ft. Spray to only wet the tall fescue blades.
Tall Fescue	Out of	Bahiagrass	Corsair 75DF up to 5.3 oz/acre. Spot treat using a hand-help sprayer delivering 1 gal of spray solution per 1000 sq. ft. Spray to only wet the tall fescue blades.
Tall Fescue	Out of	Bermudagrass	Kerb 50W (2-3 lb/acre) Or Corsair 75DF up to 5.3 oz/acre Or Manor or Blade 60DF (metsulfuron) (0.25-1 oz/acre) Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.
Tall Fescue	Out of	Zoysiagrass (Meyer & Emerald)	Manor or Blade 60DF (metsulfuron) (0.25-1 oz/acre) Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.
Tall Fescue	Out of	St. Augustine	Manor or Blade 60DF (metsulfuron) (0.25-1 oz/acre) Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.
Tall Fescue	Out of	Fine Fescue	Corsair 75DF up to 5.3 oz/acre. Spot treat using a hand-help sprayer delivering 1 gal of spray solution per 1000 sq. ft. Spray to only wet the tall fescue blades. Or Manor or Blade 60DF (metsulfuron) (0.25-1 oz/acre) Note the low use rate. As weeds mature, the rate must be increased. A nonionic surfactant at 0.25 % by volume (1qt/100 gal) increases control. Do not use beneath desirable trees or ornamentals or on desirable 'Pensacola' bahiagrass. Do not apply to desirable tall fescue or ryegrass. Some bahiagrass varieties ('Common,' 'Argentine,' & 'Paraguayan') are not completely susceptible.

*Provided by L.B. McCarty, Clemson University. — Indicates no recommendations.

Please contact your
Harrell's Representative
for more information



720 Kraft Road
Lakeland, FL 33815
800.282.8007
www.harrells.com